

SPE

Solid Phase
Extraction

Application Note

World Leaders in Sample Preparation

SPE120

ISOLATION OF NADOLOL FROM SERUM

Spe-ed™ Cartridge	Cat. No. 2001- Octadecyl C18/14%, 100mg/1mL. Process with <i>Spe-ed Mate</i> .
Sample Preparation	Dilute 1mL of sample with 1mL of 0.1 M sodium acetate.
Cartridge Conditioning	2 x 1mL of methanol, followed immediately by 1mL of 0.1 M sodium acetate. DO NOT ALLOW CARTRIDGE TO RUN DRY!
Sample Addition	Add up to 2mL of prepared sample. Aspirate sample completely from cartridge.
Cartridge Wash	2 sequential 0.25mL volumes of 70% distilled water/30% methanol, followed by 0.2mL of hexane. Air-dry column under vacuum for 2minutes.
Analyte Elution	2 sequential 0.25mL volumes of methanol. It may be necessary to evaporate sample and reconstitute in an analytically suitable solvent/ solution.

Note: *Since sample matrix interferences and concentrations may vary from sample to sample, it may be necessary to adjust the wash and elution solvent/solvent solution and/or volume to optimize isolation.*

Applied
Separations

930 Hamilton Street · Allentown, PA 18101
610-770-0900 · 610-740-5520 (fax)
www.appliedseparations.com